

Current Approaches to the Determination of Feed Intake and Digestibility in Ruminant Animals – A Review

Kubkomawa¹, H. I., Nafarnda², D. W., Adamu³, S. M., Tizhe⁴, M. A., Daniel⁵, T. K., Shua⁶, N. J., Ugwu⁷, C. C., Opara⁸, M. N., Neils⁹, J. S. and Okoli¹⁰, I. C.

1, 3, 4, 5, 6 Lecturers in the Department of Animal Production and Health, The Federal Polytechnic, P. M. B. 35, Mubi, Adamawa State, NIGERIA

2 Lecturer in the Faculty of Veterinary Medicine University of Abuja, NIGERIA

7, 8, 10 Lecturers in the Department of Animal Science and Technology, Federal University of Technology, Owerri, P. M. B. 1526, Imo State, NIGERIA

9 Lecturer in the Department of Animal Production, Adamawa State University, P.M.B. 25 Mubi Adamawa State, NIGERIA

ibrahimkubkomawa@yahoo.com

ABSTRACT

The objective of this paper is to review the current approaches for determining feed intake and digestibility in ruminant animals. Plant species or parts can differ markedly in nutritive value. The botanical composition of consumed herbage can have a profound effect on the provision of nutrients to the animal. To evaluate the preferred diet intake of grazing animals, it is necessary to spatially separate the forages being evaluated to eliminate the constraints that occur within mixed forages. In livestock production context, this is especially relevant for mixed grass and legume forages, because the consumption of legume will usually result in better animal performance. For rangeland cattle, the higher consumption of some plant species rather than others not only has nutritional effects on the animal but can also have an important influence on the species composition of the plant biomass, with consequences for ecological sustainability. Initially, quantitative estimation of the different ingested plant fractions was commonly practiced. It however, during the last four decades, many empirical models have been developed to estimate forage intake at pasture. Most of these models were derived from dairy cows experimental data, which were often specific to the relatively short range of grazing and experimental conditions used to develop them. Recent research approaches for segregating mixed swards however employ the N-alkanes profiles determination method which have been developed from different methodologies and have been used successfully to determine the proportion of different plant species in the diet. Furthermore, the oral administration of N-alkanes has been used in digestibility trials with domestic and wild ruminants as well as monogastric animals to measure feed intake and digestibility of the available herbage. Similarly, indigestible internal plant markers such as lignin and acid detergent fibres are being used to determine digestibility in ruminants. N-alkanes can be supplied to animals in different forms. Pellet feed made of paper strip embedded with synthetic N-alkanes as external markers has been used to estimate feed intake of sheep. Since evaluation of preferred forage will continue to be important in grazing lands development, researchers in regions trying to move from traditional extensive pastoral practices to some form of semi-sedentary production under managed fields may find these approaches very useful. As these novel approaches continue to evolve, it is expected that they will become simplified and cost effective and therefore find wider application in agriculture and diagnostic research.

Keywords: Forage, Feed Intake, Digestibility, N-alkane, Plant markers, Ruminants

INTRODUCTION

Forage is the most economical ruminant feed during the grazing season [1]. The establishment of mixed pasture systems by including legumes and forbs, however, has gained interest as many seek to increase biodiversity [2]. In such diverse landscapes, the ability to determine dry matter intake (DMI) and digestibility is a valuable area of study due to its impact on the nutritional status, productivity and health of animals. It also adds to our knowledge of ruminant foraging behavior and impact of seasons on biodiversity and the dynamics of the plant community [3, 4]. A considerable amount of research has been conducted using long chain saturated hydrocarbons (N-alkanes) as markers to estimate feed intake and digestibility [5, 6, 7]. N-alkanes are saturated, aliphatic hydrocarbons with length varying from 21 to 37 carbon atoms [8]. They are part of the cuticular wax of plant leaves and are usually part of the ether extract which are indigestible in nature. The oral administration of N-alkanes has been used in digestibility trials with domestic and wild ruminants as well as monogastric animals to measure feed digestibility [9, 10] and feed intake [11, 12].

N-alkanes can be supplied to animals in different forms. Pellet feed made of paper strip embedded with synthetic N-alkanes as external markers was used to estimate feed intake of sheep. Similarly, [11] fed sheep with N-alkanes (C₂₈ and C₃₂) in the form of gelatin capsules of powdered cellulose, previously added with a known amount of N-alkane dissolved with N-hexane or N-heptane, to estimate forage intake. [13] developed a different method that consists of mixing N-alkanes with solvents and powdered cellulose, resulting in a homogenous suspension that, after being evaporated and dried, was inserted into gelatin capsules. [14] developed another technique in which particles of *Pennisetum clandestinum* (Hochst.) were mixed with N-alkanes suspended in a xanthan gum (0.4%) and infused into the rumen of sheep using either drench guns or disposable syringes. In the same vein, indigestible internal plant markers such as lignin and acid detergent fibres are also used to determine feed digestibility [15, 16, 17]. Grazing time alone cannot be used to determine DMI of grazing animals because intake rate also must be considered [18]. The objective of this review is to highlight the technologies used for determination of feed intake and digestibility in ruminant animals.

Using Plant Wax Alkanes to Estimate Feed Intake and Digestibility

The history of the use of plant wax components in the study of ruminant nutrition was presented in detail by [5]. Although the use of long chain fatty acids of plant wax, (C_{29} - C_{32}), as digestibility markers was advocated. Alkanes were first examined as digestibility markers [19]. They demonstrated that in sheep, the fecal recovery of alkanes was incomplete and progressively declined as carbon-chain length decreased. This has now been confirmed in several studies reviewed in detail [5]. [20] established that negligible synthesis of alkanes occurred in the ruminant digestive tract, and although alkanes were predominantly associated with panicle matter in digesta, incomplete fecal recovery was due to absorption from the small intestine. Corrections for incomplete fecal recovery would be required if the natural alkanes were to be used on their own as markers to estimate digestibility. [19] found that the fecal recovery of C_{35} alkane was 0.975 and [5] later reported a mean C_{35} recovery in sheep of $0.948, \hat{A} \pm 0.0102$.

Based on an assumed fecal recovery of 0.95, C_{35} alkane has now been used successfully as a digestibility marker in sheep, providing estimates of digestibility which were more accurate than either in vitro estimates or those derived using lignin as a marker [21]. The validity of digestibility estimates obtained from C_{35} levels in diet and feces and an assumed fecal recovery value depends upon the degree to which the recovery value can vary. Errors in digestibility may be greater in cattle in which the fecal recovery of alkanes appears to be lower and more erratic than in sheep [5]. A further disadvantage of the use of C_{35} alkane as a marker to estimate digestibility is the fact that many plants have C_{35} in very low concentrations. [22] developed a double alkane procedure for estimating intake. In this approach, animals are dosed with known quantities of even-chain alkane and intake is estimated from the daily dose rate and the dietary and fecal concentrations of the dosed, even-chain alkane and a natural, odd-chain alkane adjacent in chain length. Note that in the feces, it is only the ratio of the alkane concentrations which is important. From their literature survey, [5] reported that alkanes C_{32} and C_{33} had recoveries of $0.868, \hat{A} \pm 0.0175$ and $0.872, \hat{A} \pm 0.0125$, respectively. Due to the similar recoveries of adjacent alkanes, comparison of known intakes with those estimated using adjacent alkane pairs show good agreement.

The response is linear and amounts to an error of 1.25% in estimated intake for every percentage unit difference in recovery between the alkane pair. As [23] and [24] have discussed, errors in the estimate of digestibility when intake is estimated from the Cr_2O_3

in- vitro procedure, result in large errors in the estimated intake, especially when digestibility is high. The oral administration of synthetic, even-chain alkanes to animals has most frequently been carried out using once- or twice-daily dosing with either pellets of alkane-impregnated shredded paper [11, 22] or gelatin capsules containing alkanes suspended on powdered cellulose [11, 13]. Procedures for preparing pellets and capsules, and for analyzing alkanes in diet and feces have been discussed elsewhere [22, 5, 25, 13, 26, 27]. The analytical method of choice involves heating the sample with 1 mol/L ethanolic KOH in a closed tube for up to 16 h, followed by extraction into N-heptane; after purification through silica gel, the hydrocarbon fraction is analyzed by capillary gas chromatography. As with dosed Cr_2O_3 , it takes 5-7 days from the commencement of alkane dosing to reach equilibrium concentrations in feces [11, 22, 16]. Within-day variation in the fecal concentration of dosed alkane is small for sheep dosed once daily with alkane-impregnated paper or twice daily with alkane on powdered cellulose, but may be higher with cattle [5]. However, the ratio of the fecal concentrations of the alkane pair used in the calculation of dietary intake is less prone to temporal variation than are the absolute concentrations [16]. To reduce the labor required for daily or more frequent dosing of animals with alkanes, which can become prohibitive in large scale intake studies, an intra-ruminal alkane CRD has been developed and its performance in sheep has been evaluated [16]. Release rates of the alkanes in the CRD were shown to be constant and were within 1.5-4.0% of the nominal release rates. The coefficient of variation of release rate between animals was 4.07%, and the intake estimated using the CRD was almost identical to known herbage intake. As with the CRD developed to administer Cr_2O_3 , the alkane CRD has advantages over the once- or twice-daily dosing, of substantially reducing disturbance to the animals and minimizing the likelihood of diurnal variation effects. In-door validation studies have thus shown that the alkane procedure for estimating dietary intake is reliable [22, 13].

However, absolute validation of the method with grazing or browsing animals is virtually impossible to achieve, because alternative methods with which to compare the technique may be no more reliable or possibly inferior. Factors which may possibly influence the reliability of the technique when used with grazing or browsing animals, such as within- and between-day variations in feeding pattern have not been extensively studied. The main precaution required in the use of the method is to ensure that the diet sample, in terms of its alkane concentrations, is representative of that consumed by the experimental animals. For uniform, sown pastures, this is relatively easy to achieve by hand-gathering or by collecting extrusa samples from fasting fistulate animals [28].

Under conditions in which animals can feed on complex vegetation communities, it may be extremely difficult or impossible to obtain feed samples having alkane concentrations which are representative of those in the diets of individual animals.

In such situations, the characterization of the botanical composition of the diet would enable intake to be assessed using the alkane technique in which the alkane concentrations of individual dietary components will have to be determined. While the alkane marker technique has been shown to give accurate assessments of dietary intake, the method has a number of other advantages. Provided that the alkane composition of the analyzed vegetation sample is representative of the diet of individual animals, the method allows for between-animal variation in diet digestibility and thus provides estimates of individual intakes. In contrast to the Cr_2O_3 in- vitro procedure [29], it can thus be used in genetic studies of the differences between individual animals in intake, digestibility and food conversion efficiency. Concurrent dosing with an external marker such as Cr_2O_3 to determine the fecal output allows the estimation of diet digestibility. However, dosed C_{36} alkane has been shown to have a consistently high fecal recovery $0.947, \hat{A} \pm 0.0139$ [5] and could thus be used in place of Cr_2O_3 . This has the advantage that fecal C_{36} concentrations can be determined in the same analysis as for the alkanes used for intake estimation. The method can accommodate the feeding of supplements to the animals, provided individual intakes of supplement are known or can be estimated by other techniques such as those discussed [21]. Supplements with negligible alkane contents can be ignored in the intake calculation. Alternatively, if the feed supplement has an alkane pattern very different from that of the basal diet, the approach for measuring the botanical composition of the diet can be adopted.

This method can also be used to obtain individual intakes in group-housed animals. Individual intakes could also be determined in large-scale trials under feedlot conditions, or in commercial feedlots in which the conversion of feed to gain is a major determinant of profitability. Whereas, alternative methods of estimating intake such as the Cr_2O_3 in -vitro technique require separate analytical procedures for fecal output and digestibility estimation, the alkane technique requires a single analytical process. As will be discussed below, the method is readily extended to the simultaneous measurement of the botanical composition of the diet, so that the intake of individual species or plant parts can be quantified.

Intake Expression

Ways of expressing level of intake are numerous; forage dry or organic matter intake is expressed in weight unit per animal per day but the expression cannot be used to compare animal species or forages between them. For this purpose, intake may be

expressed by kg of body weight raised to an exponent that can vary between 0.54 and 1.00 [30]. The choice of the exponent is a function of forage quality. With low quality forage, intake capacity of the animal appears more linked to the fill gut capacity and the rate of passage of forage. For such forages, the exponent is 1.00 and intake is expressed per kg of body weight or in percent of body weight [31]. Intake of good quality forage seems more controlled by physiological mechanisms. Intake of such forage is usually expressed per kg of metabolic weight (body weight raised to 0.75). The assumption is that intake is linked to energy requirements that are proportional to 0.75 power of body weight [32]. Nevertheless, a recent study by [33] has shown that, to compare intake level across forages and animal species, the best unit remains the dry matter intake in percent of body weight (DMI % BW). On this basis, relations between intake and passage rate of particles through the rumen or the energy digestibility appear independent of animal species. This is not the case when intake is expressed per kg of metabolic weight.

Level of Intake at Grazing

A first observation is that intake level related to animal is highly variable and linked to the characteristics of the forage. According to the French references, the *ad libitum* intake of a reference grass (15% of crude protein, 77% of organic matter digestibility on a dry matter basis) is 75, 95 and 140 g of dry matter per kg of metabolic weight respectively for a standard sheep, standard cattle (heifers) and a standard lactating dairy cow. On this basis, it is possible to calculate the "Fill Unit" ("*unité d'encombrement*") of various forages [34]. However, huge intake level variability also occurs between breed and individual animals within a given breed [35, 36]. As an example, Dorper sheep are less selective grazers as they consume more shrubs and bushes and ingest a larger number of different plant species than Merinos sheep [37].

The way of expressing intake level also contributes to this variability. As described by [33], DMI, expressed in % of body weight, appears higher for small ruminants (sheep and goats) than for cattle (dairy or suckling cattle), but this is the reverse when intake is expressed in terms of metabolic weight. Compared to temperate forages, voluntary intake of tropical forages is often lower but not highly different (2.03 kg DM per % BW vs 1.95 kg DM per % BW respectively for temperate and tropical forages), as confirmed in the meta-analysis of [38]. Despite, on a chemical point of view, temperate forages contain often higher proteins and lesser fibers. The difference decreases when the crude protein contents of both forages are similar. To explain the few differences, [38] indicated that tropical forages are usually longer chewed than temperate ones leading to a more efficient reduction of forage particles size compensating their lower apparent nutritive value.

Factors Affecting Intake Regulation

The control of intake is multi-factorial as shown by [39, 40]. It depends on plants characteristics in relation to the gut capacity, animal's requirements and nutrients concentration of forage, post-ingestive feedback of the intake and the learning process. It also depends on morphological characteristics of grazed plants, and on the environment such as climate, abundance and frequency of feed resources [41].

Role of the Ruminal Fill

The fill gut capacity, in relation to forage characteristics, is considered as a main factor of regulation of voluntary intake. Intake appears limited by the maximal volume that the digestive tract can hold [32, 42], even if herbivores modify the volume of their rumen to increase the transit rate of digesta when the quality of forage decreased [43, 44, 45]. This has been confirmed by the introduction of tennis balls, water filled bags, or artificial fibres into the rumen. The more bulky the ruminal ballast is, in volume or in weight, the more the intake decreases with or without digestibility modification [45]. In a recent study, [46] confirmed that, ruminal fill can affect grazing behavior in terms of bite mass, bite depth and bite area. In this way, it is the short-term intake that is affected by the ruminal fill. In link to the ruminal fill, forages dry matter content can influence the voluntary intake. If dry matter of forage is lower than 20%, as in young grazed grass, the volume of water in the rumen increases and has depressive effect on the intake level, this in spite of a high forage digestibility [47, 30].

The age of plant regrowth is also a factor of variation. When plant protein content decreases, the cell walls and tissue lignification increases, as a consequence, there would be an increase in forage retention time in the rumen, limiting voluntary intake [48, 41, 49]. According to [50], the daily herbage intake of lactating dairy cows decreases by 8.4% when compared with short and high age of grass regrowth. [48] and [51] confirmed that the level of dry matter intake was negatively correlated to the hemi-cellulose and cellulose (NDF) content. But as described by the low coefficient of correlation, ($r = -0.65$ and $r = -0.31$), NDF alone appears as a bad predictor of intake.

Nutrient Concentrations and Requirements

The nutrients concentrations of forages are factors of interest in the regulation of food intake. On a "requirement theory" basis, the animal eats in order to maximize its production potential under some constraints such as its gut volume and diet quality [52]. According to this theory, intake regulation would be based on the meeting of energy needs [53, 54]. [55] and [56] have demonstrated that intake is positively linked to the body weight and to the level of production of dairy cows and therefore, linked to the animal's requirements. In a meta-analysis performed

[51], factors linked to dairy cow needs and performances, such as animal body weight, change in body weight and milk yield, explained 71% of the total variations observed in dry matter intake. [57] confirmed that dry matter intake is strongly correlated to both nutrient digestibility and animal's requirements. As a consequence, ruminants eating very fibrous forages are generally unable to cover their energy needs [48].

In the area of energy needs, animal physiological state appears to be an important factor controlling voluntary intake. As an example, lactating dairy cows with higher energy requirements, graze more selectively and intensively than dry cows [58]. [43] and [44] reported that, there exist a critical rumen fill above which dry matter intake is limited. This critical rumen fill could be differently related to the physiological status of the animal. So, the introduction of ballast into the rumen of dairy cows when the energy requirements are high (beginning of lactation), involves an important decrease of intake level (0.099 to 0.043 kg dry matter per liter of added bulk). To explain energy need, cows in early lactation compensate with increasing the ruminal volume, decreasing the volume of digesta and increasing the passage rate of digesta. If the same experiment is repeated with dairy cows later in their lactation, when their energy requirements are lower, intake does not seem to be modified by the introduction of the ballast in the rumen. The pregnancy state is also a factor controlling the voluntary intake. In the predictive equation [59], intake capacity of dairy cows is proportional to lactation state, age and maturity of the cow, and to a pregnancy indicator that explains the decrease of intake during the last weeks of pregnancy. Animals can also regulate their own intake in relation to forage nutrient concentration. [60] has shown that, if sheep have the choice between two diets with different energy levels, they choose the diet that presents the highest energy density. If sheep have no choice, they adapt their intake until their requirements are met. The energy and protein balance of the diet can also influence the level of intake and the diet selection. So, lambs that select between pairs of diet, varying between 7.8 and 23.5% of crude protein, present a maximal level of intake with the diets containing between 14.1 and 17.2% of crude proteins [61]. Based on these observations, a judicious forage supplementation could contribute to improve diet nutritional balance and also increase the total voluntary intake [62, 63, 51].

Pre - and Post-Ingestive Feedback

The level of intake can be conditioned by other characteristics of the forage, such as flavor (taste and smell), appearance and texture and also by the post-ingestive feedback occurring after its intake. In other terms: "if forage tastes good, animals tend to eat it more" [64]. The flavor – feedback interaction depends

directly on chemical characteristics of feed, animal's nutritional status and animal's past or recent experiences. According to [65], the training of the animals can be made in various ways. Animals learn from their mothers before and after birth, from pair groups, testing new food and accepting or rejecting them according to the consequences induced by intake. This can explain why the preference is never fixed.

The post-ingestive feedback can evolve in relation to new experiences performed by the animal [66]. Ruminants are able to learn about the toxicity of a feed. For grazing and browsing ruminants, numerous plants contain secondary metabolites like tannins and substances that could affect digestibility, cause negative effects and reduce voluntary feed intake. Ruminants are able to use this information to modulate or avoid intake of such plants, if necessary. For instance, browsing goats tend to reduce their preference for browses sprayed with lithium chloride, chemical components associated with high negative post-ingestive feedback; tannins may reduce grazing of some forage legumes such as *Lotus pedunculatus* by affecting ruminal fermentations [67]; terpenes can inhibit cellulolytic activity of rumen micro-organisms and so limit the intake of such plants.

Finally, regulation of intake appears mediated by different signals, under the form of metabolites and hormones, emitted by the central nervous system and the peripheral organs like liver, pancreas and intestinal tracts that can be regarded as mediators of appetite [39]. The role of leptin, (hormone secreted by fatty cells), cholecystokinin, (hormone secreted by the intestinal mucous membrane), and insulin that controls satiety in ruminants have been widely documented [68, 40]. During digestion process, the rumen environment, (P^H and osmolality) can also explain the variation of voluntary intake [59, 56]. As demonstrated by [60], sheep are able to make short-term changes in diet selection to maintain good ruminal fermentations and wellbeing sensation.

Plant Morphological Characteristics

Sward characteristics, in terms of blade morphology, such as hair occurrence, thickness of cuticle [69], leaf size, stem physical properties, dead materials ratio, can stimulate, limit or inhibit animal foraging behavior [65]. These parameters have huge influence on bite size and intake rate. [70] reported that, in relation to grass characteristics, bite size can vary from 10 to 400 mg for sheep and from 70 to 610 mg for cattle. Under grazing, there is a close relationship between leaf proportion [50, 71], green leaf mass [72, 73], sward density [70] and dry matter intake.

According to [74], stems can have a barrier effect on bite size and instantaneous intake rate. The higher the stems density, the smaller the bite area and the slower the biting rate. This leads to a decrease in the instantaneous intake rate. [75] confirmed that stem

length and proportion in the sward have a negative impact on biting rate with correlation of - 0.67 and - 0.40 respectively. Sward composition, in terms of plant species, can also influence the level of intake. Indeed, compared to grasses, legumes such as white clover are often associated to higher level of intake [76]. [64, 38] explained that forage legumes are faster reduced in small particles than grasses and that less time is needed to take and masticate a similar bite of clover than grass.

Role of Environment in Feed Intake

Intake during grazing does not depend only on diet quality. Short-term intake rate is also directly correlated to forage distribution and availability [77]. This explains the lower level of intake observed under tropical rangeland, where forage resources are scattered and/or heterogeneous, which results to reduction in biting frequency and intake rate due to the time spent to go from one favorite site to another [78]. Nevertheless, animals compensate biting rate decrease by increasing grazing time. According to [58], when sward availability is measured through sward height decreases, cows increase their total grazing times, total jaw movement and total number of bites in order to maintain daily intake. These observations are confirmed on tropical forages [75].

Climatic conditions also play an important role. Ruminant animals graze essentially during daylight. In temperate climate, they make 6 to 8 meals with 2 being main meals, one at the beginning and one at the end of the day. If temperature is higher than 25°C, they adapt grazing to early morning hours or late evenings in order to avoid the hot periods of the day and this decreases the time spent grazing and total daily intake [41].

Environment also plays an important role in resource utilization. On one hand, animals have a memory of food allowance, location and distribution as related in the review of [79]. On the other hand, the rearing practices can explain the ability of cows to graze specific environment such as mountain slopes depending on the animal breeds [80]. Interaction between animals in the herd is sometimes cited to explain difference in animal grazing behavior. [81] reports that, on homogeneous vegetation, total time spent grazing by Scottish blackface sheep is higher when space allowance is high (200m² per head vs 50m² per head) without impact on herbage intake or digestibility. They conclude that the relation between time spent grazing and space allowance can be used to explain the extra activity required to maintain the group cohesion when space allowance increases.

Intake Measurement under Grazing

Direct intake measurement methods are based on herbage mass measurement. In most cases, intake is estimated by the method of difference as used [82].

The method implies the knowledge of herbage mass before and after grazing. The herbage mass is usually estimated by cutting and weighing the grass harvested on a defined area. A sward height meter or rising plate meter or disk meter may also be used to estimate grass density and quantity. The different methods are easy to apply and give reliable results if grazing period is short, (one or two days at the maximum), and stocking rate is high, (ideally all the grass of the grazing area must be consumed). If grazing period is longer, the error of estimation, linked to the grass regrowth during this period, is the major disadvantage of these methods.

To reflect the effect of grass regrowth, herbage mass and regrowth are measured in cages that exclude grazing animals. By successive cuttings, the grazing is simulated and the herbage mass accumulation is measured. But without urinary and dung restitution, specific defoliation linked to the grazing, the measured grass accumulation is often very different in grazed or non-grazed area [83]. The precision of the cutting methods is essentially based on the sampling methodology and a good precision is required at all steps of the protocol to avoid the addition of errors in the measurements. These different techniques are used to measure the intake of animal herds [82]. Instantaneous intake can also be directly estimated through live weight differences [84]. With that method, it is possible to measure intake only over a very short period (1 hour as an example). The accuracy of the measurements is strongly dependent on the precision of balance and weight loss related to dung and urine excretion during the period of measurements.

Another method of intake estimation is based on the hypothesis that the knowledge of animal requirements and performances is a good reflection of the nutritive values of ingested diet. The method is often used to determine the potential of intake of dairy cows as described [56]. Grazing supplemented dairy cows to determine grass intake from animal performances is reliable and less expensive than other methods.

Beef cattle were used to estimate intake from live weight and rate of growth with a good accuracy (residual standard deviation of 8.7% of the mean). The difficulty of the method is precisely the determination of the true herbivore requirements. This is particularly true under tropical rangeland where many external factors like displacements, feed researches must be added to basal animal needs [32].

Intake of grazing ruminants can be estimated by indirect methods such as the markers techniques, ratio techniques, the recording of animal behavior and other empirical models. The markers technique implies the determination of natural indigestible plant components, such as lignin, alkanes, or insoluble ashes which are

excreted in feces. The N-alkanes method appears as the best to estimate intake under grazing. The method, based on the determination of the concentration in plant and feces, natural alkanes and synthetic alkanes, allows the researcher to calculate intake from: $I = (F_i/F_j) \times D_j / (H_i - (F_i/F_j) \times H_j)$, where I = intake; F_i and F_j = concentration of natural and synthetic alkanes in feces; D_j = dose rate of synthetic alkanes; H_i and H_j = concentration of natural and synthetic alkanes in forage. Ratio technique implies the determination of two parameters: forage digestibility and fecal output that allows estimating intake: If $D = 100 \times (I - F) / I$, then $I = F / (1 - D/100)$ where D = forage digestibility coefficient (%); I = intake (weight unit per day); F = total fecal excretion (weight unit per day) [63].

[85] reported several methods for the determination of fecal output. The total collection of feces is one of the methods, though it is difficult to apply at grazing. To collect feces, animal must be harnessed with fecal bag or tethered, which could interrupt grazing behavior [63]. Quantity of feces that are difficult to estimate could escape from the collection bags and this could be a source of error in the estimation of intake [85]. Another method to estimate fecal output is the use of indigestible external markers as chromium oxide, ytterbium. Total collection of feces using dosing of markers requires a daily manipulation of the animals. The developed controlled release device technique, which pulses automatically a daily amount of external markers in rumen throughout the trial period, allows limiting the animal manipulation and seems sufficiently accurate to give a good estimate of fecal output [86]. Markers technique has difficulty in the sampling of forage that must be representative of the ingested diet. The N-alkanes in plant organs and plant species present different N-alkanes profiles [87] and this is a major source of error in the estimation of intake. If digestibility is determined on a non-representative sample of grazed grass, the estimation of intake will be biased. The hand plucking method, that simulates the biting of the herbivore, may be used to sample grazed grass. The measurement is linked to the calibration between animals and operator's observations. Such calibration appears easier to set up with cattle than with sheep and goats that are more selective in grazing behavior [88].

The use of oesophageal fistulate appears unfavourable to animal welfare and can modify herbivore behavior, as reported in several studies [89, 90]. Intake can be indirectly estimated by studying the grazing behavior. Indeed, intake is the product of three parameters: grazing time, biting rate and bite mass. Grazing time and biting rate can be measured by visual observation [2]. The method is easier to apply and does not require costly equipment. Nevertheless, the presence of the observer can disturb animal grazing, so, it is primordial to accustom animals to the observer in order to avoid

any behavior modification [91]. The recording of animal activities such as displacement, rumination and intake times have been largely tested and used to determine grazing time and biting rate [92]. These recording methodologies require expensive materials and the harnessing of the animal with recording apparatus can disturb its behavior. Such techniques are difficult to apply with wild herbivores and on heterogeneous rangeland.

Another alternative is the micro-histological analysis of plant residue contents in feces or the stomach and intestinal tract. The method is often used to approach the intake of wild ruminants. The main disadvantages of this technique are that, except for feces collection, it requires the slaughter of the animal and that the identification of the ingested plant fraction, till the species level, is very difficult due to the digestion process [93].

Moreover, the quantitative estimation of the different ingested plant fractions are very few as the quantity of plant fragments found in feces or in stomachs are not, due to differential digestibility, directly proportional to the quantity of the ingested plant fractions.

During the last 30 years, many empirical models have been developed to estimate forage intake at pasture. These models are based on multiple regressions between intake level and some characteristics of plant (OM yield, fiber content, digestibility, part of legumes, etc.), animal (live weight, average daily gain, milk production, stage of lactation, milk composition, pregnancy, etc.) and environment (temperature, rainfall, etc.). Most of these models derive from dairy cows experimental data [94]. They are often specific to the relatively short range of grazing and experimental conditions used to develop them.

CONCLUSION

Since evaluation of preferred forage will continue to be important in grazing lands development, it is necessary to spatially separate the forages being evaluated to eliminate the constraints that occur within an intimately mixed sward. This is most critical for researchers in regions which are trying to move from traditional extensive pastoral practices to some form of semi-sedentary production of ruminants under a managed field. As these reviewed novel approaches continue to evolve, it is expected that they will become simplified and cost effective and, therefore, find wider applications in agriculture and diagnostic research.

REFERENCES

[1] Virginie, D., André, B., Didier, S. (2009). Factors Affecting Intake by Grazing Ruminants and Related Quantification Methods: A Review *Biotechnol. Agron. Soc. Environ.* 2009, **13** (4): 559-573.

[2] Rook, A.J., Dumont, B., Isselstein, J., Osors, K., Wallis De Vries, M. F. Parnte, G., and J. Mills. (2004). Matching Type of Grazing Animal to Desired Biodiversity Outcomes—A Review. *Biol. Conserv.*, 119:137–150.

[3] Kelman, W., Bugalho, M., and Dove, H. (2003). Cuticular Wax Alkanes and Alcohols Used as Markers to Estimate Diet Composition of Sheep. *Biochem. Syst. Ecol.*, 31:919–927.

[4] Ali, H.A.M., Mayes, R.W., Lamb, C.S., Hector, B.L., Verma, A.K. and Ørskov, E.R. (2004). The Potential of Long-Chain Fatty Alcohols and Long-Chain Fatty Acids as Diet Composition Markers: Development of Methods for Quantitative Analysis and Fecal Recoveries of These Compounds in Sheep Fed Mixed Diets. *J. Agric. Sci.* 142:71–78.

[5] Dove, H. and Mayes, R. W. (1991). The Use of Plant Wax Alkanes as Marker Substance in Studies of Nutrition Herbivores: A Review. *Aust. J. Agric. Res.* **42**: 913-952.

[6] Mayes, R.W., Dove, H., Chen, X. B., and Guada, J. A. (1995). Advances in the Use of Fecal and Urinary Markers for Measuring Diet Composition, Herbage Intake and Nutrient Use in Herbivores. In: M. Journet, Editor, Recent Developments in the Nutrition of Herbivores. Proceedings of the 4th International Symposium on the Nutrition of Herbivores, Clermont-Ferrand, France. 11–15 Sept. 1995. Institut National De La Recherche Agronomique, Paris, France. P. 381–406.

[7] Dove, H. and Mayes, R.W. (2006). Protocol for the Analysis of N-Alkanes and other Plant-Wax Compounds and for their Use as Markers for Quantifying the Nutrient Supply of Large Mammalian Herbivores. *Nat. Protoc.* 1:1680–1697.

[8] Premaratne, S., Fontenot, J. P. and Shanklin, R. K. (2005). Use of N-alkanes to Estimate Intake and Digestibility by Beef Steers. *Asian-Aust. J. Anim. Sci.*, **18** (11): 1564-1568.

[9] Hulbert, I. (1993). The Effects of Upland Afforestation on the Ecology of Lagomorphs with Different Feeding Strategies. Doctoral Thesis, University of Edinburgh, Edinburgh, Scotland.

[10] Oliveira, D. E., Prates, E. R., Peralba, M. C. R. (2000). Digestibilidade Determinada Pelo Método Indireto Usando O N-Alcano C35. *Revista Brasileira De Zootecnia* 29: 848-852.

[11] Dove, H., Mayes, R. W., Freer, M., Coombe, J. B. and Foot, J. Z. (1989). Faecal Recoveries of the Alkanes of Plant Cuticular Waxes in Penned and in Grazing Sheep. *Proc. XVI Int. Grassland Cong.* Pp. 1093-1094.

[12] Oliveira, D.E.; Medeiros, S.R.; Tedeschi, L.O.; Aroeira, L.J.M.; Da Silva, S.C. (2007). Estimating Forage Intake of Lactating Dual-Purpose Cows Using Chromium Oxide and N-Alkanes as External Markers. *Scientia Agricola*, 64:103-110.

[13] Vulich, S. A., O'riordan, E. G. and Hanrahan, J. P. (1991). Use of N-Alkanes for the Estimation of Herbage

- Intake in Sheep: Accuracy and Precision of the Estimates. *J. Agric. Sci. Camb.*, 116: 319-323.
- [14] Marais, J.P.; Figenschou, P.L.; Escott-Watson, P.L.; Webber, L.N. (1996). Administration in Suspension-Form of N-Alkane External Markers for Dry Matter Intake and Diet Selection Studies. *Journal of Agricultural Science*, **126**:207-210.
- [15] Sunvold, G. D. and Cochran, R. C. (1991). Technical Note: Evaluation of Acid Detergent Lignin, Alkaline Peroxide Lignin, Acid Insoluble Ash, and Indigestible Acid Detergent Fiber as Internal Markers for Prediction of Alfalfa, Bromegrass, and Prairie Hay Digestibility by Beef Steers. *J. Anim. Sci.*, **69**: 4951-4955.
- [16] Dove, H., Mayes, R. W., Lamb, C. S. and Ellis, K. J. (1991). Evaluation of an Intra-Ruminal Controlled-Release Device For Estimating Herbage Intake Using Synthetic and Plant Cuticular Wax Alkanes. Proc. 3rd Int. Symp. Nutr. Herbivores, Pp. 82.
- [17] Fulford, G. (1994). The Potential Use of Plant Hydrocarbons Additional to N-Alkanes as Markers for Estimating Diet Composition of Large Herbivores. M.Sc. Thesis, University of Aberdeen, Aberdeen, Scotland.
- [18] Rutter, S.M. (2006). Diet Preference for Grass and Legumes in Free ranging Domestic Sheep and Cattle: Current Theory and Future Application. *Appl. Anim. Behav. Sci.*, 97:17-35.
- [19] Mayes, R. W. and Lamb, C. S. (1984). The Possible Use of N-Alkanes in Herbage as Indigestible Faecal Markers. *Proc. Nutr. Soc.*, 43: 39.
- [20] Mayes, R.W.; Lamb, C.S.; Colgrove, P.M. (1988). Digestion and Metabolism of Dosed Even-Chain and Odd-Chain N-Alkanes in Sheep. In: General Meeting European Grassland Federation, 12., Dublin, 1988. **Proceedings**. Dublin: Bsas., 159-163.
- [21] Dove, H. and Coombe, J. B. (1992). A Comparison of Methods for Estimating Supplement Intake and Diet Digestibility in Sheep. *Proc. Aust. Soc. Anim. Prod.* 19: 239-241.
- [22] Mayes, R. W., Lamb, C. S. and Colgrove, P. M. (1986a). The Use of Dosed and Herbage N-Alkanes as Markers for the Determination of Herbage Intake. *J. Agric. Sci. Camb.*, **107**: 161-170.
- [23] Langlands, J. P. (1975). Techniques for Estimating Nutrient Intake and its Utilisation by the Grazing Ruminant. In: Digestion and Metabolism in the Ruminant (McDonald, I. W. and Warner, A.C.I., Eds.), Pp. 320-332. University of New England, Armidale, Australia.
- [24] Langlands, J. P. (1987). Assessing the Nutrient Status of Ruminants. In: the Nutrition of Herbivores (Hacker, J. B. and Ternouth, J. H., Eds.), Pp. 363-390. Academic Press, Sydney, Australia.
- [25] Laredo, M. A., Simpson, G. D., Minson, D. J. and Orpin, C. G. (1991). The Potential for Using N-Alkanes in Tropical Forages as a Marker for the Determination of Dry Matter Intake by Grazing Ruminants. *J. Agric. Sci., Camb.*, **117**: 355-361.
- [26] Salt, C. A., Mayes, R. W., Colgrove, P. M. and Lamb, C. S. (1994). The Effects of Season and Diet Composition on the Radiocaesium Intake by Sheep Grazing on Heather Moorland. *J. Appl. Ecol.*, **31**:125-136.
- [27] Vulich, S. A., Hanrahan, J. P. and Crowley, B. A. (1995). Modification of the Analytical Procedures for the Determination of Herbage and Faecal N-Alkanes Used in the Estimation of Herbage Intake. *J. Agric. Sci. Camb.*, **124**: 71-77.
- [28] Vulich, S. A., Hanrahan, J. P. and O'riordan, E. G. (1993). Pasture Sampling for the Estimation of Herbage Intake Using N-Alkanes: Evaluation of Alternative Sampling Procedures. *Irish J. Agric. Res.*, **32**: 1-11.
- [29] Parker, W. J., Morris, S. T., Garrick, D. J., Vincent, G. L. and McCutcheon, S. N. (1990) Intraruminal Chromium Controlled Release Capsules for Measuring Herbage Intake in Ruminants a Review. *Proc. N. Z. Soc. Anim. Prod.*, 50: 437- 442.
- [30] Meissner, H. H. and Paulsmeier, D. V. (1995). Plant Compositional Constituents Affecting Between-Plant and Animal Species Prediction of Forage Intake. *J. Anim. Sci.*, **73**: 2447- 2457.
- [31] Demment, M. W. and Van Soest, P. J. (1985). A Nutritional Explanation for Body-Size Patterns of Ruminant and Non-Ruminant Herbivores. *Am. Nat.* **125**: 641-672.
- [32] Allison, C. D. (1985). Factors Affecting Forage Intake by Range Ruminants: A Review. *J. Range Manage.* 38(4): 305-311.
- [33] Sauvant, D., Assoumaya, C., Giger-Reverdin, S. and Archimède, H. (2006). Étude Comparative Du Mode D'expression Du Niveau D'alimentation Chez Les Ruminants. In: *Actes Des 13e Journées Autour Des Recherches Sur Les Ruminants, Rencontres Recherches Ruminants, 6-7.12.2006, Paris, France*. Paris : INRA Éditions, 103.
- [34] INRA. (2007). *Alimentation Des Bovins, Ovins Et Caprins. Besoins Des Animaux, Valeurs Des Aliments*. Versailles, France: Éditions Quæ.
- [35] Scott, L. L. and Provenza, F. D. (1999). Variation in Food Selection Among Lambs: Effects of Basal Diet and Foods Offered in a Meal. *J. Anim. Sci.*, **77**: 2391-2397.
- [36] Pearson, R. A., Archibald, R. F. and Muirhead, R. H. (2005). A Comparison of the Effect of Forage Type and Level of Feeding on the Digestibility and Gastrointestinal Mean Retention Time of Dry Forages Given to Cattle, Sheep, Ponies and Donkeys. *Br. J. Nutr.*, **95** (1): 88-98.
- [37] Brand, T. S. (2000). Grazing Behaviour and Diet Selection by Dorper Sheep. *Small Ruminant Res.* **36**(2): 147-158.
- [38] Assoumaya, C., Sauvant, D. and Archimède, H. (2007). Étude Comparative De L'ingestion Et De La Digestion Des Fourrages Tropicaux Et Tempérés. *INRA Prod. Anim.* 20 (5): 383-392.

- [39] Rhind, S. M., Archer, Z. A. and Adam, C. L. (2002). Seasonality of Food Intake in Ruminants: Recent Developments in Understanding. *Nutr. Res. Rev.*, **15**: 43-65.
- [40] Forbes, J. M. (2003). The Multifactorial Nature of Food Intake Control. *J. Anim. Sci.* **81** (2): 139-144.
- [41] Baumont, R., Prache, S., Meuret, M. and Morhand, Fehr P. (2000). How Forage Characteristics Influence Behavior and Intake in Small Ruminants: A Review. *Livestock Prod. Sci.* **61**(1): 15-28.
- [42] Allen, M. S. (1996). Physical Constraints on Voluntary Intake of Forages by Ruminants. *J. Anim. Sci.* **74**: 3063-3075.
- [43] Johnson, T. R. and Combs, D. K. (1991). Effects of Prepartum Diet, Inert Rumen Bulk, and Dietary Polyethylene Glycol on Dry Matter Intake of Lactating Dairy Cows. *J. Dairy Sci.*, **74**: 933-944.
- [44] Johnson, T. R. and Combs, D. K. (1992). Effects of Inert Rumen Bulk on Dry Matter Intake in Early and Midlactation Cows Fed Diets Differing in Forage Content. *J. Dairy Sci.*, **75**: 508-519.
- [45] Schettini, M. A., Prigge, E. C. and Nestor, E. L. (1999). Influence of Mass and Volume of Ruminant Contents on Voluntary Intake and Digesta Passage of Forage Diet in Steers. *J. Anim. Sci.*, **77**: 1896-1904.
- [46] Gregorini, P., Gunter, S. A., Masino, C. A. and Beck, P. A. (2007). Effect of Ruminant Fill on Short-Term Intake Rate and Grazing Dynamics of Beef Heifers. *Grass Forage Sci.*, **62**: 346-354.
- [47] Pasha, T. N., Prigge, E. C., Russell, R.W. and Bryan, W. B. (1994). Influence of Moisture Content of Forage Diets on Intake and Digestion by Sheep. *J. Anim. Sci.* **72** (9): 2455-2463.
- [48] Jung, H. G. and Allen, M. S. (1995). Characteristics of Plant Cell Walls Affecting Intake and Digestibility of Forages by Ruminants. *J. Anim. Sci.*, **73**: 2774-2790.
- [49] Arthington, J. D. and Brown, W. F. (2005). Estimation of Feeding Value of Four Tropical Forage Species at Two Stages of Maturity. *J. Anim. Sci.* **83**: 1726-1731.
- [50] Parga, J., Peyraud, J. L. and Delagarde, R. (2002). Age of Regrowth Affects Grass Intake and Ruminant Fermentations in Grazing Dairy Cows. In: Durand, J. L., Emile, J. C., Huyghe, C. and Lemaire G., Eds. *Proceedings of the 19th General Meeting of the European Grassland Federation on Multi-Function Grasslands, Quality Forages, Animal Products and Landscapes, 29.05.2002, La Rochelle, France*. British Grassland Society, Reading, UK: 256- 257.
- [51] Vazquez, O. P. and Smith, T. R. (2000). Factors Affecting Pasture Intake and Total Dry Matter Intake in Grazing Dairy Cows. *J. Dairy Sci.*, **83**: 2301-2309.
- [52] Yearsley, J., Tolkamp, B. J. N. and Illius, A. W. (2001). Theoretical Developments in the Study and Prediction of Food Intake. *Proc. Nutr. Soc.*, **60**.
- [53] Van Wieren, S. E. (1996). Do Large Herbivores Select a Diet that Maximizes Short-Term Energy Intake Rate? *Forest Ecol. Manage.*, **88**: 149-156.
- [54] Kyriazakis, I. (2003). What Are Ruminant Herbivores Trying to Achieve Through their Feeding Behaviour and Food Intake. In: Herrera-Camacho C.A. & Sandoval-Castro J., Eds. *Proceedings Of The 6th International Symposium On The Nutrition Of Herbivores, 9-24 October 2003, Universidad Autonoma De Yucatan, Merida, Mexico*, 153-173.
- [55] Peyraud, J. L., Comeron, E. A., Wade, M. H. and Lemaire, G. (1996). The Effect of Daily Herbage Allowance, Herbage Mass and Animal Factors Upon Herbage Intake by Grazing Dairy Cows. *Ann. Zootech.*, **45** (3): 201-217.
- [56] Faverdin, P., Delaby, L. and Delagarde, R. (2007). L'ingestion d'aliments par les vaches laitières et sa prévision au cours de la lactation. *INRA Prod. Anim.*, **20** (2): 151-162.
- [57] Hristov, A. N., Price, W. J. and Shafii, B. (2005). A Meta-Analysis on the Relationship between Intake of Nutrients and Body Weight with Milk Volume and Milk Protein Yield in Dairy Cows. *J. Dairy Sci.*, **88**: 2860-2869.
- [58] Gibb, M. J., Huckle, C. A., Nuthall, R. and Rook, A. J. (1999). The Effect of Physiological State (Lactating or Dry) and Sward Surface Height on Grazing Behaviour and Intake by Dairy Cows. *Appl. Anim. Behav. Sci.*, **63** (4): 269-287.
- [59] Faverdin, P., Delaby, L. and Delagarde, R. (2006). Prévision de l'ingestion des vaches laitières au cours de la lactation. In: *Actes des 13^e Journées autour des Recherches sur les Ruminants, Rencontres Recherches Ruminants, 6-7.12.2006, Paris, France*. Paris : INRA Éditions, 85-88.
- [60] Cooper, D. B. and Kyriazaki, I. (1995). Diet Selection in Sheep: The Role of the Rumen Environment in the Selection of a Diet From Two Feeds that Differ in their Energy Density. *Br. J. Nutr.* **74**: 39-54.
- [61] Kyriazakis, I. and Oldham, J. D. (1993). Diet selection in sheep: the ability of growing lambs to select a diet that meets their crude protein (nitrogen x 6.25) requirements. *Br. J. Nutr.*, **69**: 617-629.
- [62] Berzaghi, P., Herbein, J. H. and Polan, C. E. (1996). Intake, Site, and Extent of Nutrient Digestion of Lactating Cows Grazing Pasture. *J. Dairy Sci.* **79**(9): 1581-1589.
- [63] Lippke, H., Forbes, T. D. A. and Ellis, W. C. (2000). Effect of Supplement On Growth And Forage Intake By Stocker Steers Grazing Wheat Pasture. *J. Anim. Sci.*, **78** (6): 1625-1635.
- [64] Baumont, R. (1996). Palatabilité et Comportement Alimentaire chez le Ruminant. *INRA Prod. Anim.* **9** (5): 349-358.

- [65] Provenza, F. D. (2003a). *Foraging Behavior: Managing to Survive in a World of Change*. Washington, DC: USDA.
- [66] Atwood, S. B., Provenza, F. D., Wiedmeier, R. D. and Banner, R. E. (2001). Influence of Free-Choice Vs Mixed-Ration Diets on Food Intake and Performance of Fattening Calves. *J. Anim. Sci.* **79**: 3034-3040.
- [67] Reed, J. D. (1995). Nutritional Toxicology of Tannins and Related Polyphenols in Forage Legumes. *J. Anim. Sci.*, **73**: 1516-1528.
- [68] Forbes, J. M. (1996). Integration of Regulatory Signals Controlling Forage Intake in Ruminants. *J. Anim. Sci.*, **74**: 3029-3035.
- [69] Loney, P. E., Mc Arthur, Y. C., Potts, B. M. and Jordan G. J. (2006). How Does Ontogeny in a *Eucalyptus* Species Affect Patterns of Herbivory by Brushtail Possums? *Funct. Ecol.*, **20** (6): 982-988.
- [70] Prache, S. and Peyraud, J. L. (1997). Préhensibilité De L'herbe Pâturée Chez Les Bovins Et Les Ovins. *INRA Prod. Anim.*, **10** (5): 377-390.
- [71] O'Donovan, M. and Delaby, L. (2005). A Comparison of Perennial Ryegrass Cultivars Differing in Heading Date and Grass Ploidy With Spring Calving Dairy Cows Grazed at Two Different Stocking Rate. *Anim. Res.*, **54**: 337-350.
- [72] Penning, P. D., Parsons, A. J., Orr, R. J. and Hooper, G. E. (1994). Intake and Behaviour Responses by Sheep to Changes in Sward Characteristics under Rotational Grazing. *Grass Forage Sci.*, **49**: 476-486.
- [73] Smit, H. J., Taweel, H. Z., Tas, B. M. and Elgersma, A. (2005b). Comparison of Techniques for Estimating Herbage Intake of Grazing Dairy Cows. *J. Dairy Sci.*, **88**: 1827-1836.
- [74] Benvenuti, M. A., Gordon, I. J. and Poppi, D. P. (2006). The Effect of the Density and Physical Properties of Grass Stems on the Foraging Behaviour and Instantaneous Intake Rate by Cattle Grazing an Artificial Reproductive Tropical Sward. *Grass Forage Sci.* **61**: 272-281.
- [75] Boval, M., Fanchone, A., Archimède, H. and Gibb, M. J. (2007). Effect of Structure of a Tropical Pasture on Ingestive Behaviour, Digestibility of Diet and Daily Intake by Grazing Cattle. *Grass Forage Sci.* **62**: 44-54.
- [76] Ribeiro, H. M. N., Delagarde, R. and Peyraud, J. L. (2003). Inclusion of White Clover in Strip-Grazed Ryegrass Swards: Herbage Intake and Milk Yield of Dairy Cows at Different Ages of Sward Regrowth. *Anim. Sci.*, **77**: 499-510.
- [77] Garcia, F., Carrère, P., Soussana, J. F. and Baumont, R. (2003). How do Severity and Frequency of Grazing Affect Sward Characteristics and the Choices of Sheep during the Grazing Season? *Grass Forage Sci.*, **58**: 138-150.
- [78] Roguet, C., Dumont, B. and Prache, S. (1998). Sélection Et Utilisation Des Ressources Fourragères Par Les
- [79] Baker, S. K. and Klein, L. (1994). Potential to Use N-Alkanes in Plant Cuticular Waxes to Discriminate Plant Parts of Subterranean Clovers Eaten by Ruminants. *Proc. Aust. Soc. Anim. Prod.* **20**: 419.
- [80] Tulloch, A. P., Baum, B. R. and Hoffman, L. L. (1980). A Survey of Epicuticular Waxes among Genera of Triticeae. 2. Chemistry. *Can. J. Bot.*, **58**: 2602 - 2615.
- [81] Sibbald, A. M., Shellard, L. J. and Smart, T. S. (2000). Effect of Space Allowance on the Grazing Behaviour and Spacing of Sheep. *Appl. Anim. Behav. Sci.* **70** (1): 49-62.
- [82] Newman, J. A., Thompson, W. A., Penning, P. D. and Mayes, R. W. (1995). Least-Squares Estimation of Diet Composition-From Nalkanes in Herbage and Faeces Using Matrix Mathematics. *Aust. J. Agric. Res.*, **46**: 793 - 805.
- [83] Frame, J. (1993). Herbage Mass. In: Davies A., Baker R.D., Grand S.A. & Laidlaw A.S., eds. *Handbook*. 2nd ed. Hurley, Berkshire, UK: British Grassland Society, 39-68.
- [84] Coates, D. B. and Penning, P. (2000). Measuring Animal Performances. In: 'T Mannetje L. and Jones R.M., Eds. *Field and Laboratory Methods for Grassland and Animal Production Research*. Wallingford, UK: CABI Publishing, 353-402.
- [85] Adegoosan, A. T., Givens, D. I. and Owens, E. (2000). Measuring Chemical Composition and Nutritive Values in Forages. *Field and Laboratory Methods for Grassland and Animal Production Research*. Wallingford, UK: CAB International, 263-278.
- [86] Compère, R. (1992). Comparaison De Deux Methodes Utilisant Le Cr2O3 Pour Estimer La Quantité De Feces Émises Par Les Moutons : Pellets De Luzerne Et Bolus CAPTEC-CRD. *Bull. Rech. Agron. Gembloux*, **27**: 365-383.
- [87] Cortes, C., Damanesco, J. C., Bechet, G. and Prache, S. (2005). Species Composition of Ryegrass (*Lolium Perenne*) and Tall Fescue (*Festuca Arundinacea*) Mixtures Using Various Combinations of N-Alkanes. *Grass Forage Sci.* **60**: 254-261.
- [88] DeVries, W. M. F. (1995). Estimating Forage Intake and Quality in Grazing Cattle: A Reconsideration of the Handplucking Method. *J. Range Manage.* **48**: 370-375.
- [89] Coates, D. B., Schachenmann, P. and Jones, R. J. (1987). Reliability of Extrusa Samples Collected from Steers Fistulated at the Oesophagus to Estimate the Diet of Resident Animals in Grazing Experiments. *Aust. J. Exp. Agric.* **27**: 739-745.
- [90] Jones, R. J. and Lascano, C. E. (1992). Oesophageal Fistulated Cattle Can Give Unreliable Estimates of the Proportion of Legume in the Diets of Resident Animals Grazing Tropical Pastures. *Grass Forage Sci.*, **47**: 128-132.

- [91] Agreil, C. and Meuret, M. (2004). An Improved Method for Quantifying Intake Rate and Ingestive Behaviour of Ruminants in Diverse and Variable Habitats Using Direct Observation. *Small Ruminant Res.* 54 (1-2): 99-113.
- [92] Laca, E. A. and Wallis de Vries, M. F. (2000). Acoustic Measurement of Intake and Grazing Behaviour of Cattle. *Grass Forage Sci.*, **55**: 97-104.
- [93] Holechek, J. L., Vavra, M. and Piepper, R. D. (1982a). Botanical Composition Determination of Range Herbivores Diets: A Review. *J. Range Manage.*, **35** (3): 309 - 315.
- [94] Holter, J. B., West, J. W. and Mc Gilliard, M. L. (1997). Predicting Ad-Libitum Dry Matter Intake and Yield of Holstein Cows. *J. Dairy Sci.*, **80**: 2188-2199.